

POSSIBILITIES OF BASIC INHALATIVE THERAPY IN ASTHMA PATIENTS

*U. I. Feshchenko, L. O. Iashyna,
T. O. Pertseva, M. O. Polianska, T. V. Kireeva*

Summary

Own experience of 2 months course of combined medicine Airtec (fluticasone propionate|salmeterol) in two doses: 125/25 μg (in steroid-naïve patients and in patients with not adequate control on low ICS doses in monotherapy) and 250/25 μg – in patients with not adequate control on moderate ICS doses or low doses of ICS/LABA. Use of this medicine allowed to improve quality of therapy – not to lose control in patients on combined ICS/LABA therapy, to improve asthma control in adults with persistiv BA, steroid-naïve, those who received ICS in monotherapy or with nonadequate previous therapy with low dosed of ICS/LABA.

Key words: *bronchial asthma, control, combined therapy.*

Theoretical and practical J. «Asthma and Allergy», 2015, 3

Y. I. Feshchenko

Academician of NAMS of Ukraine, professor

*SI «National Institute of phthisiology and pulmonology
named after F. G. Yanovskii NAMS of Ukraine»*

Amosova str., 10, Kyiv, 03680

tel.: +38(044)275-04-02; +38(044)275-21-18

e-mail: admin@ifp.kiev.ua